

Representation theorem for obseables on a quantum logic

Andrei Khrenikov

International Centre for Mathematical Modeling
in Physics and Cognitive Sciences,
MSI, University of Växjö, S-35195 Sweden
E-mail: Andrei.Khrennikov@msi.vxu.se

Olga Nánásiová

Department of Mathematics and Descriptive Geometry
Faculty of Civil Engineering
Slovak University of Technology
Radlinského 11, 813 68 Bratislava, Slovakia
Email: olga@res.svf.stuba.sk

February 1, 2008

Abstract

We will study a conditional state on a quantum logic using Renyi's approach (or Bayesian principle). This approach helps us to define independence of events and differently from the situation in the classical theory of probability, if an event a is independent of an event b , then the event b can be dependent on the event a . We will show that we can define a s -map (function for simultaneous measurements on a quantum logic). It can be shown that if we have the conditional state we can define the s -map and conversely. By using the s -map we can introduce joint distribution also for noncompatible observables on a quantum logic.

1 Introduction

The idea that quantum probabilities are nothing other than conditional probabilities was intensively discussed from various points of views, see, e.g., [1]–[5]. Recently this approach to quantum probability was generalized in the so called contextual framework [6]–[11]. In the latter approach we consider conditioning with respect to various contexts, complexes of experimental physical conditions and not conditioning of one event, a , with respect to other event, b , under the same context (state) as it was done in investigations on conditional probabilities (compare with Kolmogorov [12], [13], Renyi [14], Cox [15], or quantum logic generalizations [16], see also [1]–[5]).

In the present paper we still use the traditional event-conditioning (under the fixed conditional state). However, we essentially generalize the notion of a conditional state (here we continue investigations [18]–[23]).

We will study a conditional state on a quantum logic using Renyi's approach (or Bayesian principle). This approach helps us to define independence of events and differently from the situation in the classical theory of probability, if an event a is independent of an event b , then the event b can be dependent on the event a (problem of causality) ([22], [23]). We will show that we can define a s -map (function for simultaneous measurements on a quantum logic). It can be shown that if we have the conditional state we can define the s -map and conversely. By using the s -map we can introduce joint distribution also for noncompatible observables on a quantum logic. Moreover, if x is an observable on L and B is Boolean sub-algebra of L , we can construct an observable $z = Exp(x|B)$, which is a version of conditional expectation of x but it need not to be necessarily compatible with x .

1. A conditional state on a quantum logic

In this part we introduce the notions as a quantum logic, a state, a conditional state and their basic properties.

Definition 1.1 *Let L be a nonempty set endowed with a partial ordering $\leq$. Let there exists the greatest element (1) and the smallest element (0). Let there be defined the operations supremum ($\vee$), infimum ($\wedge$) (the lattice operations) and a map $\perp: L \rightarrow L$ with the following properties:*

(i) *For any $\{a_n\}_{n \in \mathcal{A}} \in L$, where $\mathcal{A} \subset \mathcal{N}$ are finite*

$$\bigvee_{n \in \mathcal{A}} a_n, \bigwedge_{n \in \mathcal{A}} a_n \in L.$$

(ii) *For any $a \in L$ $(a^\perp)^\perp = a$.*

(iii) *If $a \in L$, then $a \vee a^\perp = 1$.*

(iv) *If $a, b \in L$ such that $a \leq b$, then $b^\perp \leq a^\perp$.*

(v) *If $a, b \in L$ such that $a \leq b$ then $b = a \vee (a^\perp \wedge b)$ (orthomodular law).*

Then $(L, 0, 1, \vee, \wedge, \perp)$ is called the quantum logic (briefly q.l.).

Let L be q.l.. Then elements $a, b \in L$ will be called:

- *orthogonal* ($a \perp b$) iff $a \leq b^\perp$;

- *compatible* ($a \leftrightarrow b$) iff there exist mutually orthogonal elements $a_1, b_1, c \in L$ such that

$$a = a_1 \vee c \quad \text{and} \quad b = b_1 \vee c.$$

If $a_i \in L$ for any $i = 1, 2, 3, \dots$ and $b \in L$ is such, that $b \leftrightarrow a_i$ for all i , then $b \leftrightarrow \bigvee_{i=1}^n a_i$ and

$$b \wedge \left(\bigvee_{i=1}^{\infty} a_i \right) = \bigvee_{i=1}^{\infty} (a_i \wedge b)$$

([17],[24],[26]).

A subset $L_0 \subseteq L$ is a *sub-logic* of L if for any $a \in L_0$ we have $a^\perp \in L_0$ and for any $a, b \in L_0$ $a \vee b \in L_0$.

Definition 1. 2 A map $m : L \rightarrow [0, \infty)$ such that

(i) $m(0) = 0$ and $m(1) = 1$.

(ii) If $a \perp b$ then $m(a \vee b) = m(a) + m(b)$

is called a *state* on L . If we have orthomodular σ -lattice and m is σ -additive function, then m will be called a σ -state.

Definition 1. 3 [23] Let L be a quantum logic. A subset $L_c \subset L - \{0\}$ is called a *conditional system (CS)* in L (σ -CS in L) if the following conditions hold:

- If $a, b \in L_c$, then $a \vee b \in L_c$. (If $a_n \in L_c$, for $n = 1, 2, \dots$, then $\bigvee_n a_n \in L_c$.)
- If $a, b \in L_c$ and $a < b$, then $a^\perp \wedge b \in L_c$.

Let $A \subset L$. Then $L_c(A)$ is the smallest CS (σ -CS), which contains the set A .

Definition 1. 4 [23] Let L be a quantum logic and L_c be an σ -CS in L . Let $f : L \times L_c \rightarrow [0, 1]$. If the function f fulfill the following conditions:

(C1) for each $a \in L_0$ $f(\cdot, a)$ is a state on L ;

(C2) for each $a \in L_0$ $f(a, a) = 1$;

(C3) if $\{a_n\}_{n \in \mathcal{A}} \in L_0$, where $\mathcal{A} \subset \mathcal{N}$ and a_n are mutually orthogonal, then for each $b \in L$

$$f(b, \bigvee_{n \in \mathcal{A}} a_n) = \sum_{n \in \mathcal{A}} f(a_n, \bigvee_{n \in \mathcal{A}} a_n) f(b, a_n);$$

then it is called *conditional state*.

Proposition 1. 1 [23] Let L be a q.l.. Let $\{a_i\}_{i=1}^n \in L$, $n \in \mathbb{N}$ where $a_i \perp a_j$ for $i \neq j$. If for any i there exists a state α_i , such that $\alpha_i(a_i) = 1$, then there exists σ -CS such that for any $\mathbf{k} = (k_1, k_2, \dots, k_n)$, where $k_i \in [0; 1]$ for $i \in \{1, 2, \dots, n\}$ with the property $\sum_{i=1}^n k_i = 1$, there exists a conditional state

$$f_{\mathbf{k}} : L \times L_c \rightarrow [0; 1],$$

such that

1. for any i and each $d \in L$ $f_{\mathbf{k}}(d, a_i) = \alpha_i(d)$;
2. for each a_i

$$f_{\mathbf{k}}(a_i, \bigvee_{i=1}^n a_i) = k_i;$$

Definition 1. 5 [23] Let L be a quantum logic and f be a conditional state. Let $b \in L$, $a, c \in L_c$ such that $f(c, a) = 1$. Then b is independent of a with respect to the state $f(\cdot, c)$ ($b \asymp_{f(\cdot, c)} a$) if $f(b, c) = f(b, a)$.

The classical definition of independency of a probability space $(\Omega, \mathcal{B}, \mathcal{P})$ is a special case of this definition, because

$$P(A|B) = P(A|\Omega) \text{ if and only if } P(A \cap B|\Omega) = P(A|\Omega)P(B|\Omega).$$

If L_c be CS and $f : L \times L_c \rightarrow [0, 1]$ is a conditional state, then ([23])

- (i) Let $a^\perp, a, c \in L_c$, $b \in L$ and $f(c, a) = f(c, a^\perp) = 1$. Then $b \asymp_{f(\cdot, c)} a$ if and only if $b \asymp_{f(\cdot, c)} a^\perp$.
- (ii) Let $a, c \in L_c$, $b \in L$ and $f(c, a) = 1$. Then $b \asymp_{f(\cdot, c)} a$ if and only if $b^\perp \asymp_{f(\cdot, c)} a$.
- (iii) Let $a, c, b \in L_c$, $b \leftrightarrow a$ and $f(c, a) = f(c, b) = 1$. Then $b \asymp_{f(\cdot, c)} a$ if and only if $a \asymp_{f(\cdot, c)} b$.

2. Observables and s -map

Let L be a quantum logics. Let us denote $L^2 = L \times L$.

Definition 2. 1 Let L be a quantum logic. The map $p : L^2 \rightarrow [0, 1]$ will be called s -map if the following conditions hold:

- (s1) $p(1, 1) = 1$;
- (s2) if there exists $i \in N$ $a \perp b$, then $p(a, b) = 0$;
- (s3) if $a \perp b$, then for any $c \in L$,

$$p(a \vee b, c) = p(a, c) + p(b, c)$$

$$p(c, a \vee b) = p(c, a) + p(c, b)$$

Proposition 2. 1 *Let L be a q.l. and let p be a s-map. Let $a, b, c \in L$, then*

- 1. if $a \leftrightarrow b$, then $p(a, b) = p(a \wedge b, a \wedge b) = p(b, a)$;
- 2. if $a \leq b$, then $p(a, b) = p(a, a)$;
- 3. if $a \leq b$, then $p(a, c) \leq p(b, c)$;
- 4. $p(a, b) \leq p(b, b)$;
- 5. if $\nu(b) = p(b, b)$, then ν is a state on L .

Proposition 2. 2 *Let L be a q.l., let there be a s-map p . Then there exists a conditional state f_p , such that*

$$p(a, b) = f_p(a, b)f_p(a, 1).$$

Let L be a q.l. and let $L_c = L - \{0\}$. If $f : L \times L_c \rightarrow [0, 1]$ is a conditional state, then there exists a s-map $p_f : L \times L \rightarrow [0, 1]$.

Proposition 2. 3 *Let L be a quantum logic.*

- (a) *If f is a conditional state, then $b \succ_{f(.,1)} a$ iff $p_f(b, a) = p_f(a, a)p_f(b, b)$, where p_f is the s-map generated by f .*
- (b) *Let p be a s-map. Then $b \succ_{p(.,1)} a$ iff $p(b, a) = p(a, a)p(b, b)$, where f_p is the conditional state generated by the s-map p .*

Let $\mathcal{B}(\mathcal{R})$ be σ -algebra of Borel sets. A σ -homomorphism $x : \mathcal{B}(\mathcal{R}) \rightarrow \mathcal{L}$ is called an observable on L . If x is an observable, then $R(x) := \{x(E); E \in \mathcal{B}(\mathcal{R})\}$ is called range of the observable x . It is clear that $R(x)$ is Boolean σ -algebra [26]. A spectrum of an observable x is defined by the following way: $\sigma(x) = \cap\{E \in \mathcal{B}(\mathcal{R}); x(E) = 1\}$. If g is a real function, then $g \circ x$ is such observable on L that:

- (1.) $R(g \circ x) \subset R(x)$;
- (2.) $\sigma(g \circ x) = \{g(t); \quad t \in \sigma(x)\}$;
- (3.) for any $E \in \mathcal{B}(\mathcal{R})$

$$g \circ x(E) = x(\{t \in \sigma(x); g(t) \in E\}).$$

We say that x and y are compatible ($x \leftrightarrow y$) if there exist a Boolean sub- σ -algebra $\mathcal{B} \subset L$ such that $R(x) \cup R(y) \subset \mathcal{B}$. In other words $x \leftrightarrow y$ if for any $E, F \in \mathcal{B}(\mathcal{R})$ $x(E) \leftrightarrow y(F)$.

We call an observable x a discrete if $\sigma(x)$ is a countable set. It means, that $\sigma(x) = \{t_i\}_{i \in \mathcal{A}}$, where $\mathcal{A} \subset \{1, 2, \dots\}$. Let us denote $\mathcal{O}$ the set of all discrete observables on L .

Definition 2. 2 Let L be a σ -quantum logic and $p : L \times L \rightarrow [0; 1]$ be a s -map. Let x, y be some observables on L . Then a map $p_{x,y} : \mathcal{B}(\mathcal{R}) \times \mathcal{B}(\mathcal{R}) \rightarrow [0, 1]$, such that

$$p_{x,y}(E, F) = p(x(E), y(F)),$$

is called a joint distribution for the observables x and y for the s -map p .

If x is an observable on L and m is a state on L , then $m_x(E) = m(x(E))$, $E \in \mathcal{B}(\mathcal{R})$ is a probability distribution for x and

$$m(x) = \int_R \lambda m_x(d\lambda)$$

is called the expectation of x in the state m , if the integral on the right side exists. From this follows that for $x \in \mathcal{O}$

$$m(x) = \sum_{t \in \sigma(x)} t m(x(\{t\}))$$

and for any real function g we have

$$m(g \circ x) := \sum_{t \in \sigma(x)} g(t) m(x(\{t\})).$$

Definition 2. 3 Let L be a σ -quantum logic and $p : L \times L \rightarrow [0; 1]$ be a s -map. Let $x, y \in \mathcal{O}$. Then

$$p(x, y) = \sum_{x_i \in \sigma(x)} \sum_{y_j \in \sigma(y)} x_i y_j p(x(\{x_i\}), y(\{y_j\})),$$

is called the first joint moment for the observables x, y .

From the previous definition follows that

$$p(x, x) = \sum_{x_i \in \sigma(x)} x_i^2 p(x(\{x_i\}), x(\{x_i\})) = \sum_{x_i \in \sigma(x)} x_i^2 \nu(x(\{x_i\})) := \nu(g \circ x),$$

where $g(t) = t^2$.

Analogically to the classical theory of probability we can define notions for example as covariance ($c(.,.)$), variance ($var(.)$) and correlation coefficient ($r(.,.)$) by the following way:

$$c(x, y) := p(x, y) - \nu(x)\nu(y),$$

$$var(x) := c(x, x),$$

$$r(x, y) = \frac{c(x, y)}{\sqrt{var(x).var(y)}}.$$

In spite of the classical theory of probability in this case $c(x, y)$ is not equal to $c(y, x)$ in general.

Proposition 2. 4 *Let L be a quantum logic, p be a s-map on L and $\mathcal{O}$ be a set of all discrete observables on L . For each $x, y \in \mathcal{O}$ there exist $(\Omega_i, \mathcal{S}_i, P_i)$ ($i = 1, 2$) probability spaces and random variables ξ_i, η_i ($i = 1, 2$), which are $\mathcal{S}_i$ -measurable such that:*

- (a) $c(x, y) = cov(\xi_1, \eta_1)$, $c(y, x) = cov(\eta_2, \xi_2)$;
- (b) $E_i(\xi_i) = \nu(x)$ and $E_i(\eta_i) = \nu(y)$, $i = 1, 2$;
- (c) $(c(x, y))^2 \leq c(x, x)c(y, y)$.

Proof. If $x, y \in \mathcal{O}$, then $\sigma(x) = \{x_k\}_{k \in \mathcal{A}}$, $\sigma(y) = \{y_j\}_{j \in \mathcal{D}}$, where $\mathcal{A}, \mathcal{D} \subset N$. Let us denote $\Omega_1 = \{(x_k, y_j); k \in \mathcal{A}, j \in \mathcal{D}\}$, $\Omega_2 = \{(y_j, x_k); k \in \mathcal{A}, j \in \mathcal{D}\}$ and $\mathcal{S}_i = 2^{\Omega_i}$, $i = 1, 2$. Then $(\Omega_i, \mathcal{S}_i)$ is the measurable space.

If p is a s-map, then from the properties of p follows, that $P_1 = p_{x,y}$ is the probability measure on $(\Omega_1, \mathcal{S}_1)$ and $P_2 = p_{y,x}$ is the probability measure on the measurable space $(\Omega_2, \mathcal{S}_2)$. From this

$$E_i(\xi_i) = \sum_k x_k P_i(\xi_1 = x_k) = \nu(x).$$

Analogically we get

$$E_i(\eta_i) = \nu(y).$$

(b) Let us denote

$$\xi_1((x_k, y_j)) = x_k, \quad \eta_1((x_k, y_j)) = y_j,$$

$$\xi_2((y_j, x_k)) = x_k, \quad \eta_2((y_j, x_k)) = y_j.$$

It is clear, that

$$P_1(\{\omega \in \Omega_1; \quad \xi_1(\omega) = x_k\}) = P_1\{(x_k, y_j); j \in \mathcal{D}\}$$

From this follows, that

$$P_1(\{\omega \in \Omega_1; \quad \xi_1(\omega) = x_k\}) = p(x(\{x_k\}), y(\sigma(y))) = p(x(\{x_k\}), 1) = \nu(x(\{x_k\})).$$

Anallogicaly from this follows, that

$$P_i(\xi_i = x_k) = \nu(x(\{x_k\}))$$

and

$$P_i(\eta_i = y_j) = \nu(y(\{y_j\}))$$

(a) From the theory of probability follows, that

$$var(\xi_1, \eta_1) = \sum_k \sum_j (x_k - \nu(x))(y_j - \nu(y))P_1(\xi_1 = x_k, \eta_1 = y_j))$$

Let us denote $a_k = x(\{x_k\})$ and $b_j = y(\{y_j\})$. Then

$$cov(\xi_1, \eta_1) = \sum_k (x_k - \nu(x))(y_j - \nu(y))p(a_k, b_j)$$

and

$$cov(\xi_2, \eta_2) = \sum_k (x_k - \nu(x))(y_j - \nu(y))p(b_j, a_k).$$

Because

$$\begin{aligned} \sum_{k,j} p(a_k, b_j) &= \sum_{k,j} p(b_j, a_k) = 1, \\ \sum_k p(a_k, b_j) &= \sum_k p(b_j, a_k) = \nu(b_j) \end{aligned}$$

and

$$\sum_j p(a_k, b_j) = \sum_j p(b_j, a_k) = \nu(a_k)$$

we have

$$\sum_k (x_k - \nu(x))(y_j - \nu(y))p(b_j, a_k) = p(x, y) - \nu(x)\nu(y) = c(x, y).$$

Anallogically

$$var(\xi_2, \eta_2) = p(y, x) - \nu(x)\nu(y) = c(y, x).$$

(c) Because $(cov(\xi_i, \eta_i))^2 \leq cov(\xi_i, \xi_i)cov(\eta_i, \eta_i)$ and $cov(\xi_i, \xi_i) = c(x, x)$, $cov(\eta_i, \eta_i) = c(y, y)$ we have

$$(c(x, y))^2 \leq c(x, x)c(y, y),$$

$$(c(y, x))^2 \leq c(x, x)c(y, y).$$

(Q.E.D.)

Proposition 2. 5 *Let L be a quantum logic and $\mathcal{O}$ be the set of all discrete observables on L . Let $x, y \in \mathcal{O}$. Then*

(i) $c(x, y) = p(g_x \circ x, g_y \circ y)$, where g_z is a real function such that $g_z(t) = t - \nu(z)$, for $z \in \mathcal{O}$;

(ii) $r(x, y) \in [-1, 1]$;

(iii) if $x \leftrightarrow y$, then $c(x, y) = c(y, x)$ and $r(x, y) = r(y, x)$.

Proof. Let $x, y \in \mathcal{O}$. Then $\sigma(x) = \{x_i\}_{i \in \mathcal{A}}$ and $\sigma(y) = \{y_j\}_{j \in \mathcal{B}}$, where $\mathcal{A}, \mathcal{B}$ are countable sets of indexes.

(i) From the definition $g_x \circ x, g_y \circ y$ we have

$$\begin{aligned} p(g_x \circ x, g_y \circ y) &= \sum_{x_i \in \sigma(x)} \sum_{y_j \in \sigma(y)} g_x(x_i) g_y(y_j) p(x(x_i), y(y_j)) \\ &= \sum_{x_i \in \sigma(x)} \sum_{y_j \in \sigma(y)} (x_i - \nu(x)) (y_j - \nu(y)) p(x(x_i), y(y_j)) \\ &= \sum_{x_i \in \sigma(x)} \sum_{y_j \in \sigma(y)} x_i y_j p(x(x_i), y(y_j)) \\ &\quad - \sum_{x_i \in \sigma(x)} \sum_{y_j \in \sigma(y)} \nu(x) \nu(y) p(x(x_i), y(y_j)) \\ &= p(x, y) - \nu(x) \nu(y) = c(x, y). \end{aligned}$$

(ii) From the previous proposition we know that there exist two probability spaces $(\Omega_k, \mathcal{F}_k, P_k)$ and random variables ξ_k, η_k , $k = 1, 2$ such that $c(x, y) = \text{cov}(\xi_1, \eta_1)$ and $c(x, y) = \text{cov}(\xi_2, \eta_2)$. Moreover $c(x, x) = \text{cov}(\xi_k, \xi_k) = \text{var}(\xi_k)$. Analogically $c(y, y) = \text{cov}(\eta_k, \eta_k) = \text{var}(\eta_k)$. Then

$$r(x, y) = \frac{c(x, y)}{\sqrt{c(x, x)c(y, y)}} = \frac{\text{cov}(\xi_1, \eta_1)}{\sqrt{\text{var}(\xi_1)\text{var}(\eta_1)}} = \rho(\xi_1, \eta_1) = \rho_1$$

and

$$r(y, x) = \frac{c(y, x)}{\sqrt{c(x, x)c(y, y)}} = \frac{\text{cov}(\xi_2, \eta_2)}{\sqrt{\text{var}(\xi_2)\text{var}(\eta_2)}} = \rho(\xi_2, \eta_2) = \rho_2.$$

because ρ_k is correlation coefficient on the probability space $(\Omega_k, \mathcal{F}_k, P_k)$, then $\rho_k \in [-1, 1]$ for $k = 1, 2$. From this follows that $r(x, y), r(y, x) \in [-1, 1]$. Also this fact follows immediately from the Proposition 2.4 (c).

(iii) Let $x \leftrightarrow y$. Then for each $x_i \in \sigma(x)$ and each $y_j \in \sigma(y)$ we have

$$p(x(x_i), y(y_j)) = p(y(y_j), x(x_i))$$

and so

$$\begin{aligned} p(x, y) &= \sum_{x_i \in \sigma(x)} \sum_{y_j \in \sigma(y)} x_i y_j p(x(x_i), y(y_j)) \\ &= \sum_{x_i \in \sigma(x)} \sum_{y_j \in \sigma(y)} x_i y_j p(y(y_j), x(x_i)) \\ &= p(y, x). \end{aligned}$$

From this follows that

$$c(x, y) = c(y, x).$$

(Q.E.D.)

Example 2. 1 . Let $L = \{a, a^\perp, b, b^\perp, 0, 1\}$. Let $c \vee d = 1$ if $c \neq d$ and $c, d \in L - \{0\}$. Let $c \wedge d = 0$ if $c \neq d$ and $c, d \in L - \{1\}$. Let $(d^\perp)^\perp = d$ for $d \in L$ and $1^\perp = 0$. It is clear that L is a quantum logic and $B_d = \{d, d^\perp, 0, 1\}$, $d \in \{a, b\}$ is Boolean algebra. Let $f(s, t)$ is defined by the following way:

s/t	a	$a^\perp$	b	$b^\perp$	1
a	1	0	0.4	0.4	0.4
$a^\perp$	0	1	0.6	0.6	0.6
b	0.2	11/30	1	0	0.3
$b^\perp$	0.8	19/30	0	1	0.7

From f this we can compute $p_f(s, t)$. Then we get:

s/t	a	$a^\perp$	b	$b^\perp$
a	0.4	0	0.12	0.28
$a^\perp$	0	0.6	0.18	0.42
b	0.08	0.22	0.3	0
$b^\perp$	0.32	0.38	0	0.7

We can see that $p_f(a, b) = p_f(a, a)p_f(b, b)$, but $p_f(b, a) \neq p_f(b, b)p_f(a, a)$. In the following we will write $p_f = p$. Let x, y be observales on L such that $R(x) = \{a, a^\perp, 0, 1\} = B_a$, and $R(y) = \{b, b^\perp, 0, 1\} = B_b$. It is easy to see, that x is not compatible with y . Let, for example,

$$x(-1) = a \quad x(1) = a^\perp$$

$$y(0) = b \quad y(5) = b^\perp.$$

In the following tables we have the joint distributions $p_{x,y}$ and $p_{y,x}$.

$p_{x,y}$	0	5
-1	0.12	0.28
1	0.18	0.42

$p_{y,x}$	-1	1
0	0.08	0.22
5	0.32	0.38

Now we can compute the following characteristics:

$$\nu(x) = -1 \times 0.4 + 0.2 \times 0.6 = 0.2,$$

$$\nu(y) = 0 \times 0.3 + 5 \times 0.7 = 3.5,$$

$$p(x, y) = -5 \times 0.28 + 5 \times 0.42 = 0.7,$$

$$p(y, x) = -5 \times 0.32 + 5 \times 0.38 = 0.3,$$

$$c(x, y) = p(x, y) - \nu(x)\nu(y) = 0.7 - 0.2 \times 3.5 = 0,$$

$$\begin{aligned}
c(y, x) &= p(y, x) - \nu(x)\nu(y) = 0.3 - 0.2 \times 3.5 = -0.4, \\
c(x, x) &= 0.96 & c(y, y) &= 5.25, \\
r(x, y) &= 0 & r(y, x) &\doteq 0.178.
\end{aligned}$$

In the end we can write these results to "the covariance matrix":

$$\begin{pmatrix} c(x, x) & c(x, y) \\ c(y, x) & c(y, y) \end{pmatrix} = \begin{pmatrix} 0.96 & 0 \\ 0.178 & 5.25 \end{pmatrix}$$

We see that "the covariance matrix" need not to be symmetry. In the classical theory of probability, where we suppose that all random variables are compatible, have to be symmetry.

Example 2. 2 . Let L the same quantum logic as in the Example 2.1. Let $p(s, t)$ is defined by the following way:

s/t	a	$a^\perp$	b	$b^\perp$
a	0.4	0	0.08	0.38
$a^\perp$	0	0.6	0.22	0.32
b	0.08	0.22	0.3	0
$b^\perp$	0.32	0.38	0	0.7

Let x, y be observables on L such that $R(x) = \{a, a^\perp, 0, 1\}$, and $R(y) = \{b, b^\perp, 0, 1\}$. It is easy to see, that x is not compatible with y . Let, for example,

$$\begin{aligned}
x(-1) &= a & x(1) &= a^\perp \\
y(0) &= b & y(5) &= b^\perp.
\end{aligned}$$

In this case $p_{x,y} = p_{y,x}$.

$p_{x,y}$	0	5
-1	0.08	0.38
1	0.22	0.32

Now we get

$$\begin{aligned}
\nu(x) &= 0.2, & \nu(y) &= 3.5, \\
p(x, y) &= p(y, x) = -0.3, & c(x, y) &= c(y, x) = -0.4, \\
c(x, x) &= 0.96, & c(y, y) &= 5.25, \\
r(x, y) &= r(y, x) \doteq 0.178.
\end{aligned}$$

In the end we can write these results to the covariance matrix:

$$\begin{pmatrix} c(x, x) & c(x, y) \\ c(y, x) & c(y, y) \end{pmatrix} = \begin{pmatrix} 0.96 & 0.178 \\ 0.178 & 5.25 \end{pmatrix}$$

We see that the covariance matrix is symmetry as in the classical theory of probability, but x, y are not compatible.

In the end we can say that we cannot prove that two observables are compatible by using statistics but we can prove that they are not compatible.

References

- [1] Accardi L., The probabilistic roots of the quantum mechanical paradoxes. *The wave-particle dualism. A tribute to Louis de Broglie on his 90th Birthday*, ed. S. Diner, D. Fargue, G. Lochak and F. Selleri (D. Reidel Publ. Company, Dordrecht, 297–330, 1984).
- [2] Ballentine L. E. , Probability theory in quantum mechanics. *American J. of Physics*, **54**, 883-888 (1986).
- [3] Ballentine L. E., Interpretations of probability and quantum theory. Proc. Conf. *Foundations of Probability and Physics*, ed. A. Khrennikov. *Q. Prob. White Noise Anal.*, **13**, 71-84, WSP, Singapore (2001).
- [4] Gudder S. P., An approach to quantum probability. Proc. Conf. *Foundations of Probability and Physics*, ed. A. Khrennikov. Quantum Prob. White Noise Anal., **13**, 147-160, WSP, Singapore (2001).
- [5] De Muynck W. M., Interpretations of quantum mechanics, and interpretations of violation of Bell's inequality. Proc. Conf. *Foundations of Probability and Physics*, ed. A. Khrennikov. *Q. Prob. White Noise Anal.*, **13**, 95-114, WSP, Singapore (2001).
- [6] Khrennikov A. Yu., *Contextual viewpoint to quantum stochastics*. hep-th/0112076.
- [7] Khrennikov A. Yu., *Interpretations of Probability*, VSP Int. Sc. Publ., Utrecht (1999).
- [8] Khrennikov A. Yu., Linear representations of probabilistic transformations induced by context transitions. *J. Phys.A: Math. Gen.*, **34**, 9965-9981 (2001).
- [9] Khrennikov A. Yu., *Hyperbolic quantum mechanics*. quant-ph/0101002 (2000).
- [10] Khrennikov A. Yu., Contextualist viewpoint to Greenberger-Horne-Zeilinger paradox. *Phys. Lett., A*, **278**, 307-314 (2001).
- [11] Khrennikov A. Yu., Frequency analysis of the EPR-Bell argumentation. *Foundations of Physics*, **32**, 1159-1174 (2002).
- [12] Kolmogoroff A.N., *Grundbegriffe der Wahrscheinlichkeitsrechnung*, Springer, Berlin, (1933)

- [13] Kolmogorov A. N., *The Theory of Probability*. In: A.D.Alexandrov, A.N.Kolmogorov, M.A.Lavrent'ev (Eds.) *Mathematics, Its Content, Methods, and Meaning*, **2**, M.I.T. Press (1965).
- [14] Renyi A., On a new axiomatics of probability theory. *Acta Mat. Acad. Sc. Hung.*, **6**, 285-335 (1955).
- [15] Cox R., *The algebra of probable inference*. J. Hopkins Univ. Press, Baltimore (1961).
- [16] Beltrametti E. and Cassinelli G., *The logic of quantum mechanics*. Addison-Wesley, Reading, Mass (1981).
- [17] Dvurečenskij A., Pulmannová S., *New Trends in Quantum Structures*, Kluwer Acad. Publ. (2000).
- [18] Nánásiová O., On conditional probabilities on quantum logic. *Int. J. Theor. Phys.*, **25**, 155 - 162 (1987).
- [19] Nánásiová O., Ordering of observables and characterization of conditional expectation on a quantum logic. *Math. Slovaca*, **37**, 323-340 (1987).
- [20] Nánásiová O., Observables and expectation on the Pták sum. *Tatra Mount. Publ. Math.*, **3**, 65-76 (1993).
- [21] Nánásiová O., States and homomorphism on the Pták sum. *Int. J. Theor. Pphys.*, **32**, 1957-1964 (1993).
- [22] Nánásiová O., A note on the independent events on a quantum logic. *Busefal*, **76**, 53-57 (1998).
- [23] Nánásiová O., Principle conditioning. Preprint (2001).
- [24] Pták P., Pulmannová S., *Quantum Logics*. Kluwer Acad. Press, Bratislava (1991).
- [25] Riečan B., Neubrun T., *Measure theory*.
- [26] Varadarajan V., *Geometry of quantum theory*, Princeton, New Jersey, D. Van Nostrand, (1968)